Mon. Not. R. Astron. Soc. **000**, 1–17 (2002) Printed 2 February 2008 (MN LATEX style file v2.2)

arXiv:astro-ph/0301324v2 29 Apr 2003

# Mass limits for the progenitor star of supernova 2001du and other type II-P supernovae

S. J. Smartt[1], J. R. Maund[1], G. F. Gilmore[1], C. A. Tout[1], D. Kilkenny[2], S. Benetti[3],
[1] *Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, England*
[2] *South African Astronomical Observatory, PO Box 9, Observatory 7935, South Africa*
[3] *INAF - Osservatorio Astronomico di Padova, vicolo dell'Osservatorio 5, I-35122 Padova, Italy*

Accepted. Received 2003 January; in original form 2003 January

## ABSTRACT

The supernova SN 2001du was discovered in the galaxy NGC1365 at a distance of $19 \pm 2$ Mpc, and is a core-collapse event of type II-P. Images of this galaxy, of moderate depth, have been taken with the Hubble Space Telescope approximately 6.6 years before discovery and include the supernova position on the WFPC2 field of view. We have observed the supernova with the WFPC2 to allow accurate differential astrometry of SN 2001du on the pre-explosion frames. As a core-collapse event it is expected that the progenitor was a massive, luminous star. There is a marginal detection ($3\sigma$) of a source close to the supernova position on the prediscovery $V-$band frame, however it is not precisely coincident and we do not believe it to be a robust detection of a point source. We conclude that there is no stellar progenitor at the supernova position and derive sensitivity limits of the prediscovery images which provide an upper mass limit for the progenitor star. We estimate that the progenitor had a mass of less than 15M$_\odot$. We revisit two other nearby SNe II-P which have high quality pre-explosion images, and refine the upper mass limits for the progenitor stars. Using a new distance determination for SN 1999gi from the expanding photosphere method, we revise the upper mass limit to 12M$_\odot$. We present new HST images of the site of SN 1999em, which validate the use of lower spatial resolution ground-based images in the progenitor studies and use a new Cepheid distance to the galaxy to measure an upper mass limit of 15M$_\odot$ for that progenitor. Finally we compile all the direct information available for the progenitors of eight nearby core-collapse supernovae and compare their mass estimates. These are compared with the latest stellar evolutionary models of pre-supernova evolution which have attempted to relate metallicity and mass to the supernovae type. Although this is statistically limited at present, reasonable agreement is already found for the lower mass events (generally the II-P), but some discrepancies appear at higher masses.

**Key words:** galaxies: individual (NGC1365) – stars: evolution – supernovae: general – supernovae: individual (SN 2001du, SN 1999em, SN 1999gi)

## 1 INTRODUCTION

The end point of the lives of all stars with initial mass greater than approximately 8M$_\odot$ is thought to be a core-collapse supernova. Supernovae (SNe) explosions which arise from massive stars have been a major driver in the chemical evolution of the Universe, are important in shaping the interstellar medium dynamics in gas-rich galaxies and are fundamental in the study of the origin of the chemical elements in the Universe. The exact evolutionary states of stars before they undergo core-collapse is of great interest in the fields of massive star evolution and the physics of supernovae evolution.

Supernovae are classified according to the atomic lines observed in the early-time optical spectra. Briefly, the presence of H I optical lines indicate a type II classification (SNe II), while those that show no hydrogen at all are type I (Filippenko 1997). The SNe Ia are thought to arise through thermonuclear explosions in white dwarf binary systems (Branch et al. 1995). Given the occurrence of these supernovae in both old and intermediate age stellar populations (e.g. in galaxies of type SO and earlier), the white dwarf binary scenario is consistent with the progenitors being low−intermediate mass stars. However very little other details are known about the stellar systems that produce SNe Ia. All other supernovae including the type Ib/Ic (SNe Ib/Ic) and all the subtypes of SNe II are thought to be due to core-collapse during the deaths of massive stars. The subtype SNe II-P show prominent, broad H I lines with P-Cygni type profiles in their optical spectra, indicating that the progenitor retained a substantial hydrogen envelope prior to explosion e.g. see Woosley & Weaver (1986). They also display a plateau phase in their lightcurve during which the supernovae remain at approximately constant brightness for a period of between 30 and 100 days, which is also interpreted as evidence of an extended hydrogen atmosphere in the progenitor star, such as red supergiant (Chevalier 1976, Woosley & Weaver 1986). This plateau phase is most obvious in the $V-$band, but can less pronounced or non-existant at other visual and NIR wavelengths. The type-IIn supernovae show strong hydrogen features which are in emission with broad wings and very narrow cores. These features are thought to arise from interaction with dense circumstellar material which may have been ejected from the progenitor star through stellar winds or more dramatic mass-loss events e.g. see Fassia et al. (2001), Pastorello et al. (2002), Lentz et al. (2001) and references therein. The SNe Ib/Ic also do not show any obvious signs of hydrogen in their spectra, although SNe Ib display pronounced He I absorption. However recently Branch (2003) has suggested that at least some Ib events do show some amounts of hydrogen, and some Ic events do show helium. It is likely there is a continuum of possibilities rather than the well defined sub-classes traditionally employed.

There is very strong evidence that the SNe II and Ib/Ic are associated with the deaths of massive stars as they are never seen in elliptical galaxies, only rarely in S0 types (van den Bergh et al. 2002) and they often appear to be associated with sites of recent starformation such as H II regions and OB associations in spiral and irregular galaxies (Van Dyk et al. 1996). However the desired direct evidence of detecting stars before they exploded has only been possible for a small number of objects. The two most unambiguous detections of a progenitor object of confirmed supernovae explosions are for SNe 1987A and 1993J. The progenitor to SN 1987A in the LMC (White & Malin 1987) was the blue supergiant Sk−69°202 of spectral type B3 Ia (Walborn et al. 1989). The closest supernova to the Milky Way since then was SN 1993J in M81 (3.63 Mpc), which was a type IIb event and ground based $UBVRI$ photometry of the SN site before explosion was presented by Aldering et al. (1994). The photometry of the progenitor candidate was best fit with a composite spectral energy distribution of a K0 Ia star and some excess $UB$ band flux either from unresolved OB association contamination or a hot companion. Neither the progenitor of SN 1987A nor that of SN 1993J is consistent with the canonical stellar evolution picture, where core carbon burning finishes and core-collapse occurs relatively soon afterwards ($\sim 10^3 - 10^4$ yrs) while the massive star is an M-supergiant. Claims have been made on the detection of the progenitors of three other supernovae events SNe 1961V, 1978K and 1997bs, however for reasons we discuss in Section 5.3 it is uncertain whether these were actual supernovae explosions.

The only way to further add to our knowledge of the core-collapse process and its relation to massive stellar evolution is to identify more progenitor objects. There is now a wealth of data in the Hubble Space Telescope (HST) archive, as well as high quality ground-based images from various well maintained archive facilities (e.g. ESO, ING, CFHT). Thus there are reasonable chances of having available an image of a pre-supernova site which has sufficient resolution to identify individual massive stars. Clearly there is a distance limit within which images are useful from depth and resolution constraints. This is roughly 10 Mpc for a deep, good quality (natural seeing of typically $0.7''$) image from a ground-based 2-8m telescope, and roughly 20 Mpc for typical exposure length HST frames. Recently Smartt et al. (2001a; 2002a; 2002b) have used both HST and ground based images of pre-explosion supernovae sites to try to detect progenitor stars of SNe II-P 1999gi and 1999em and the type Ic SN 2002ap. These nearby events have high quality prediscovery archive images available, but no progenitor star is actually identified in any of these cases. These papers discuss the luminosity limits which can be placed on the exploding star from the sensitivity measurements of the images, and quite restrictive mass-limits can be determined (see Section 5.3 for a full comparison and discussion). In a similar vein, Van Dyk et al. (2003a) have attempted to identify the progenitors of 16 type II and type Ib/s SNe on HST prediscovery images. However 9 of the objects in this lengthy paper are further than ∼40 Mpc, with another reddened by some $5 - 6$ magnitudes of visual extinction, hence there is little or no chance of identifying a progenitor star for these events. The other six are subject to the detailed HST follow-up programme of Smartt et al. (GO 9353) which will allow more accurate astrometry and possible source detection. The Van Dyk et al. study has attempted absolute astrometry of WFPC2 archive images as a method of determining the supernovae positions on the prediscovery frames. This method has not proved accurate enough to position the events to the precision necessary to assess if a progenitor star can be detected. The error boxes the define are generally too large to provide useful constraints. They do sugges that there are possible progenitor candidates for four supernovae, although these will require more detailed follow-up to confirm or discount the claims.

In this paper we present a search for a progenitor of the core-collapse supernova SN 2001du. This was a type II-P event in the galaxy NGC1365 at a distance of $(m - M) = 31.39 \pm 0.26$ (Ferrarese et al 2000; corresponding to $19{\pm}2$ Mpc) and the supernova position was observed by HST in 1995. The supernova was discovered by Evans (2001) on 2001 August 24.7 UT at a visual magnitude of about 14.0. It was classified by Smartt et al. (2001c) on 2001 September 2.05 UT, and Wang et al. (2001) on August 30, as a type II-P supernova due to the presence of strong H I P-Cygni profiles in the H$\beta$ and H$\alpha$ transitions. Wang et al. also obtained spectropolarimetry of SN 2001du and reported the supernova to be polarized at a level of 0.29% with a flat spectrum characteristic of electron scattering, indicative of a fairly spherical explosion. This supernova has very similar optical spectra to the other type II-P events SNe 1999em and 1999gi for which we determined upper mass limits for the progenitor star. Images of the SN 2001du host galaxy are available in the HST archive allowing another detailed search for a precursor object. In this paper we show that there is a marginal detection of a faint source close to the supernova position. But it is neither a robust detection of a single point source nor precisely coincident with the supernova. We argue that there is no detection of a progenitor, and hence determine mass and luminosity limits of the exploding star and discuss the implications of this finding alongside all the available, direct information we have on progenitors of core-collapse supernovae.

## 2 OBSERVATIONAL DATA AND ANALYSIS

### 2.1 Astrometry and detection limits of WFPC2 images

The galaxy NGC1365 was observed on four epochs with the WFPC2 on board HST before the supernova 2001du was discovered. However on only one of these occasions does the supernova position actually fall on any of the WFPC2 chips. The galaxy was a target for the HST key project on the extragalactic distance scale, which was designed to accurately resolve and photometrically measure Cepheid variables in this spiral. Unfortunately these deep exposures (totalling 18.5hrs in F555W and 6hrs in F814W) do not cover the position of the progenitor of SN 2001du. The observations which do include the supernova position were taken in Cycle 4 in GO5222 (PI: J. Trauger), and the details are listed in Table 1. These data were located and retrieved from the ST-ECF Science archive using the ASTROVIRTEL initiative. The approximate position of the supernova on these prediscovery images can be estimated (Van Dyk et al. 2003a), however one is limited by the absolute astrometric resolution of the WFPC2 frames (usually of the order $1-2''$) and the error on the SN position ($\sim 0.5-1''$). As this could amount to $\sim$30 pixels on the WF chips of WFPC2, it is essential that more accurate astrometry is achieved to determine the SN position on the prediscovery frames. As discussed in Smartt et al. (2001a), by far the best method is to take repeat WFPC2 observations of the supernova some months after explosion and use the surrounding stars in the post and pre-explosion images to determine precise *differential* astrometry. The crowded host region of SN 1999gi shown by Smartt et al. (2001a) and Li et al. (2002) illustrates that an RMS error of even $0.2''$ in positioning the supernova would not have allowed two nearby, resolved point sources to be have been ruled out as the progenitor objects. Hence re-observing the SN at the resolution of HST (with either WFPC2 or ACS) is essential to accurately determine the SN position.

The supernova was observed with WFPC2 on 2001 November 26 through the program GO9041, and the exposure times and filter details are listed in Table 1. These are total exposure times, and as the SN was still quite bright at this epoch, the F336W, F555W and F814W were broken down into sub-exposures to allow photometry of a non-saturated SN point-spread-function (PSF) as well as the fainter background stars. Both the F555W and F814W datasets are composed of a 40s and 3×200s exposures, while the F336W is composed of 100s and 460s exposures (with CR splits for cosmic ray removal). Photometry of stars visible in both these long and short exposures showed excellent alignment to better than 0.1 pixel in each dimension. Hence the centroid position of the supernova can be measured in each of the short frames, and assumed to be appropriate for the longer frames. Seventeen bright ($V_{555} \simeq 20.5-23.5$), single stars were identified in a $500 \times 500$ pixel region around the SN position in the F555W post-explosion 600s exposure and their positions measured with aperture photometry within the IRAF DAOPHOT package. The positions of these same 17 stars were identified in the prediscovery frames and a geometric solution mapping these positions to the post-explosion frame was calculated. The spatial transformation fitted a linear shift, magnification factor and a rotation, giving an RMS to the fit of $0.01''$ in each of $x$ and $y$. Eleven stars were used in the F814W transformation, with similar RMS errors. In Fig. 1 we show the position of the supernova in the 3 colour prediscovery images. The total exposure time available for all the stacked images at this position is 330s in each filter. However this is made up of 3×100s and 3×10s individual frames, Given the significant read-out noise of the WFPC2 chips (7e$^-$ per frame) compared with the sky background in a stellar aperture, the 3×10s frames were ignored, and only the 300s stacked images were used. We note that Van Dyk et al. (2003a) quote an exposure time of 100s total for these images, and presume this is a typographical error.

Very close to the position of the supernova on the F555W image there is a possible detection of a faint source which is close to the detection limit of the image. Assuming Poisson statistics, a total CCD readout noise of 12e$^-$ (from the three individual stacked frames), and a gain of 14e$^-$/ADU, this object is detected with a $3.0\sigma$ significance above the background noise. The magnitude of the object estimated by placing a 2-pixel radius aperture at the position is $V_{555} \simeq 25.0 \pm 0.4$. This was obtained using the IRAF DAOPHOT package, and applying the appropriate charge-transfer-efficiency (Whitmore et al. 1999) and aperture corrections (Holtzman et al. 1995a). A model point-spread-function was calculated using the Tiny Tim software (Krist & Hook 1999), and PSF fitting photometry was also carried out on the source using the DAOPHOT task ALLSTAR. A consistent magnitude of $V_{555} \simeq 25.2 \pm 0.4$ is measured. However it is a marginal detection, the object is not clearly identified by eye on the frames and is not a convincing point source. Placing a 2 pixel aperture centred on various pixels in the vicinity which are individually more than $2\sigma$ above their neighbours will frequently give a measured magnitude of between 25.0−26.2. The source appears somewhat extended, and has two peaks of intensity separated by roughly two pixels. After PSF subtraction one of these pixel peaks always remains, which suggests to us that this is not a robust detection of a point source. The source is not detected in the F814W filter. Furthermore, the supernova position is not exactly coincident with the object within the estimated errors of our methods. We estimate the error radius on the precursor object position to be 0.3 pixels, which is the maximum difference between positional determinations using four methods i.e. intensity weighted means in each dimension, a Gaussian fit, a triangular weighting function, and the PSF fit. However we find the object 1.03 pixels from the supernova centroid. The error in the supernova centroid is 0.14 pixels, and transformation error as discussed above is 0.14 pixels RMS, giving a total error circle of radius $r_{\rm err} = 0.28$ pixels. Hence considering the er-

**Table 1.** Details of the HST WFPC2 archival and GO observational data for SN 2001du. The dataset names come from the the ST-ECF Science archive (http://archive.eso.org) and are the associations of shorter, aligned exposures. The location column indicates where SN 2001du occurred on the WFPC2 FOV. The PC1 has a plate scale of 0.0455″ per pixel, WF3 has 0.0996″ per pixel.

| Dataset | Date | Filter | Exposure Time | Location |
|---|---|---|---|---|
| U2KV010AB | 1995 January 15th | F160BW | 330s | WF3 |
| U2KV010AB | 1995 January 15th | F336W | 330s | WF3 |
| U2KV010JB | 1995 January 15th | F555W | 330s | WF3 |
| U2KV010SB | 1995 January 15th | F814W | 330s | WF3 |
| U6BR030JB | 2001 November 26th | F336W | 560s | PC1 |
| U6BR030NB | 2001 November 26th | F439W | 400s | PC1 |
| U6BR0303B | 2001 November 26th | F555W | 640s | PC1 |
| U6BR030LB | 2001 November 26th | F675W | 400s | PC1 |
| U6BR030BB | 2001 November 26th | F814W | 640s | PC1 |

**Figure 1.** Panels (a)-(c) show images of the F336W, F555W and F814W filters taken 6.6 yrs before explosion transformed to the pixel coordinates of the post-explosion (d). The images (a)-(c) are from the WF3 chip (0.1″ pixels), and (d) is better sampled with the 0.05″ pixels of the PC1. The position of the SN is marked as a cross in the prediscovery images, and the blank triangular regions are due to the edge of the WF3 chip after transformation. The fields of view of these panels are 20″, corresponding to a distance of 1.8 kpc at the distance of NGC1365

rors in the analysis, the supernova position is not precisely coincident with this possible source (see Fig. 2).

We have followed the labelling of stars in the region from Van Dyk et al. (2003a), who identify three objects within a 0.9″ error circle of their estimated supernova position, but the only unambiguous detection we confirm is that of star A. The object we marginally detect close to the supernova position is the Van Dyk et al. star B, and we keep the nomenclature for clarity. Their star C is also a marginal detection, but as it is some way from the supernova position, we do not consider it any further. These images are available in the HST archive, and we quote our determination of the position of SN 2001du (at pixel (106.80, 675.96), with errors as listed above) on chip WF3, to allow our results to be independently verified if necessary. There doesn't appear to be a significant discrepancy between our results and those of Van Dyk et al., as the latter have used the HSTphot photometry routines of Dolphin (2000a; 2000b) with a $3\sigma$ threshold. We do repeat a $3\sigma$ detection, but for the reasons outlined above do not interpret it as a certain detection of a stellar progenitor. Van Dyk et al. measure a magnitude of 25.02±0.3, which is in good agreement with our PSF magnitude given the errors, the faintness of the source and the different methods employed.

We determined the detection limits of the data in the vicinity of the SN position using two methods. The first is just a simple calculation of the limiting sensitivities based on Poisson statistics assuming the CCD parameters given above. For a 2 pixel aperture, these are $V_{555}(3\sigma) = 25.3$ and $V_{555}(5\sigma) = 24.8$. Correcting for the CTE and applying an aperture correction would reduce these to $V_{555}(3\sigma) = 25.0$ and $V_{555}(5\sigma) = 24.5$. To validate this result, we performed simulations of a synthetic star close to the supernovae position, (at a point with very similar sky values to those surrounding the possible star B). Stellar photometry using PSF fitting procedures within the IRAF package DAOPHOT was carried out, and all the stellar sources within the region of the supernova were subtracted from the image. The Tiny Tim package (Krist & Hook 1999) was used to construct a suitable PSF for this fitting procedure. Model PSFs of varying magnitudes between 24.4−25.6 were made, again using Tiny Tim, and these were added to the "flat" image. A synthetic star at the magnitude of star B ($V_{555} \simeq$25.0) is not recovered with convincing significance. Placing an aperture at the stellar position gives a measurement of $V_{555} \simeq 25.7\pm0.9$. At the theoretical $5\sigma$ limit of $V_{555} \simeq$24.5, the object can be visually identified and a magnitude of 24.8±0.5 is determined.

We recognise that there is some statistical validity in the $3\sigma$ detection of star B by Van Dyk et al. (2003a) and ourselves. However we cannot confirm, with any certainty, an unambiguous detection of a point source. It looks more likely to us that this object is a fluctuation of a fairly complex non-stellar background, which could loosely be interpreted as a point source, and that the progenitor is below (and possibly just below) the sensitivity limit. In the rest of this paper we assume that we do not have a secure identification of the progenitor, and base our results on the robust $5\sigma$ upper limit to its $V_{555}$ magnitude. We agree with Van Dyk et al. that there is no sign of this object in the other three filters, and exactly the same method for artificial star additions was applied to F814W and F336W, resulting in similar $5\sigma$ detections at $I_{814}$=24.8 and $U_{336}$=21.6. The image in the UV filter also shows nothing to a limiting magnitude of $m_{\rm F160BW} = 19.8$, and as this isn't particularly useful for placing constraints on the progenitor or its environment it is not discussed any further.

### 2.2 Photometry of the stellar population around SN 2001du

Photometry in the three filter images F555W, F675W and F814W taken after explosion on 26 November 2001 was carried out in order to help determine a reddening towards the surrounding stellar populations. These images were chosen over the prediscovery frames as they are deeper and have



**Figure 2.** A close up of the original F555W and F814W pre-explosion images before the transformation described in Section 2.1. The position of the SN is marked at position (0,0) with the orthogonal lines. The width of these lines represent the combined error in its position and the transformation RMS fit errors (0.14″). We keep the same labeling as Van Dyk et al. (2003a) for stars A-B-C, and extend it to the other objects detected in the field. The possible object (star B) detected at $3\sigma$ above the background noise is labeled, and its position is marked with a solid dot. The size of this dot represents the probable centroid error on the position of the source, and within the errors it is not coincident with the supernova. Star B is difficult to identify visually, and is more likely a variation in the complex background rather than a true point source. It is not detected in the F814W filter. The size of the image panels are 5″, corresponding to a distance of of 460 pc at the distance of NGC1365. The directions of north and east are marked.

better spatial resolution (as SN 2001du is on the PC1 chip). Photometry using PSF fitting techniques was carried out on the stars on the PC1 chip only, as the reddening towards other parts of the galaxy contained in the rest of the WFPC2 chips is not applicable to our present study. The on-the-fly recalibrated images were taken from the archive and the IRAF version of DAOPHOT was used to do the photometry. Aperture photometry was first carried out with a 4.4 pixel radial aperture. Model PSFs were calculated using the Tiny Tim package (Krist & Hook 1999) in each filter, and these were used in the DAOPHOT task ALLSTAR. The zero-points from the WFPC2 headers were applied, and several corrections where included. Firstly the charge transfer efficiency (CTE) correction was determined (Whitmore et al. 1999), then the magnitudes were corrected for the 4.4 pixel aperture using the encircled energy tables of Holtzmann et al. (1995a). Finally a colour transformation was calculated to determine magnitudes of the stars in the Johnson $VRI$ bands from the WFPC2 flight system (Holtzman et al. 1995b). Results from the photometry are discussed below in Section 4.1.3.

## 2.3 Spectroscopy of SN 2001du

Optical spectra of SN 2001du were taken on two epochs at the South African Astronomical Observatory (SAAO), and at the European Southern Observatory. The first spectrum was taken on 2001 September 2.05 at the SAAO 1.9m using the Cassegrain Grating Spectrograph, with the No. 6 grating (600 $\mathrm{lmm}^{-1}$) and SITe detector providing a coverage of $3500-5400$Å. The spectral dispersion was 1.1Å $\mathrm{pix}^{-1}$, and the resolution measured from the width of the arclines was approximately 3Å. Our second spectrum was taken at ESO-La Silla on the Danish 1.54m telescope on 2001 September 9.35. The DFOSC spectrometer was employed with grism No. 4 and the 4k×2k EEV CCD which provided a dispersion of 2.8 Å $\mathrm{pix}^{-1}$, and useful coverage between $3300-9200$Å. The spectra were reduced using standard techniques within IRAF. The bias level was subtracted from the CCD images, before flat-fielding, extraction and wavelength calibration from arc line maps. A photometric standard star was observed on each occasion for relative flux calibration. Both of the reduced spectra are presented in Fig. 3.

Smartt et al. (2001c) and Wang et al. (2001) classified this as a type II-P supernova from the appearance of the P-cygni like H I Balmer lines, particularly H$\beta$. The unpublished lightcurve of Suntzeff & Krisciunas as shown in Van Dyk et al. (2003b) shows it to have had a plateau phase of approximately 100 days. The spectrum and lightcurve is very similar to the well studied SN 1999em which was a classic II-P event with a 95 day plateau (Hamuy et al. 2001; Leonard et al. 2002b; Elmhamdi et al. 2003), and further discussion of the similarities are discussed below in Section 4.1.1.



**Figure 3. (a):** The dereddened early time spectrum of SN 2001du taken on Sept. 2.05 is compared with black-body continuum fluxes of various temperatures (dotted lines). Comparisons with an unphysically large temperature of $T_{\rm BB} = 10^9$K effectively set a hard upper limit to the reddening. If the spectrum is dereddened by more than $E(B-V)$ >0.6, then the continuum cannot be matched by any temperature. For comparison the more likely temperatures of the photosphere of $T_{\rm BB}$ = 12000K and 20000K (see Baron et al. 2000), are shown with the reddening required to match. **(b):** Comparisons of the optical spectra of SN 2001du (heavy line) with SN 1999em (lighter line – from Hamuy et al. 2001). The upper spectra are SN 2001du taken on Sept. 2.05 and SN 1999em taken +7 days after explosion. The continuum of SN 2001du is very blue similar to SN 1999em, and indicates the reddening of the two cannot be very different. The difference in the slopes is likely due to the difference of around 4 days in the epochs. The lower spectra are SN 1999em taken on day +18 after explosion, and SN 2001du taken on 2001 September 9.35 UT. The spectra are almost identical (after a linear scaling), indicating that SN 2001du is most probably also around +18 days, and the reddening must be similar to that for SN 1999em, for which a value of $E(B-V) = 0.1 \pm 0.05$ was measured.

## 3 THEORETICAL STELLAR EVOLUTIONARY MODELS

We have calculated a number of stellar evolutionary models to compare with the luminosity constraints from the pre-discovery images. Models were constructed at every integer mass between 9−20M$_\odot$ which is the most interesting region probed by the observational data, plus several higher masses at 25, 40 and 60M$_\odot$. As described in Smartt et al. (2002a), the models come from the most recent version of the Eggleton evolution program (Eggleton 1971; Eggleton 1972; Eggleton 1973). The equation of state, which includes molecular hydrogen, pressure ionization and coulomb interactions, is discussed by Pols et al. (1995). The initial composition is taken to be uniform with a hydrogen abundance $X = 0.7$, helium $Y = 0.28$ and metals $Z = 0.02$ with the meteoritic mixture determined by Anders & Grevesse (1989). Hydrogen burning is allowed by the pp chain and the CNO cycles. Helium burning is explicitly included in the triple $\alpha$ reactions and reactions with $^{12}$C, $^{14}$N and $^{16}$O along with carbon burning via $^{12}{\rm C} +^{12}{\rm C}$ only and the disintegration of $^{20}$Ne. Other isotopes and reactions are not explicitly followed. Reaction rates are taken from Caughlan and Fowler (1988). Opacity tables are those calculated by Iglesias, Rogers and Wilson (1992) and Alexander and Ferguson (1994). An Eddington approximation (Woolley and Stibbs 1953) is used for the surface boundary conditions at an optical depth of $\tau = 2/3$. This means that low-temperature atmospheres, in which convection extends out as far as $\tau \approx 0.01$ (Baraffe et al. 1995), are not modelled perfectly. However the effect of this approximation on observable quantities is not significant in this work (see for example Kroupa and Tout 1997). We included convective overshooting in this run of models as described in Pols et al. (1997) in order to compare consistently with the models of the Geneva group which incorporate this mixing process. There is no mass-loss added to the $9-20$M$_\odot$ tracks as this plays only a very minor role in the evolution of these stars. Mass loss for the higher mass stars is included as described in Dray et al. (2003) according to their NL prescription.

The most interesting luminosity range which is of relevance to this paper, and the studies of the progenitors of SNe II-P, corresponds to the $9-20$M$_\odot$ mass range. For these masses the end points of the evolutionary tracks at the end of core-carbon burning in our tracks are very close, but not identical, to those of the Geneva group (Schaller et al. 1992;

Meynet et al. 1994). The Geneva tracks are calculated at 9, 12, 15 and 20$M_\odot$, and we have reproduced these and all integer values in between. Our tracks between 9-15$M_\odot$ end at a luminosity 0.1−0.15 dex higher than those of the Geneva group, and at 20$M_\odot$ they are virtually identical. One should note that when convective overshooting is included, then 2nd dredge-up does not occur. Smartt et al. (2002a) discussed the effects of 2nd-dredge in 7-11$M_\odot$ models without convective overshooting, which predicts that stars of this mass can achieve significantly higher theoretical luminosities. At this point however this serves as an illustrative example of the uncertainties in our understanding of the convective and mixing processes in massive stars, and the final luminosities of the 7-11$M_\odot$ stars cannot yet observationally constrain the models. This area requires more careful scrutiny from both the observational and theoretical side, and we intend to study the latter in future publications using our stellar evolutionary code. Given the fact that the pre-supernova luminosities in the 9-15$M_\odot$ range are approximately 0.15 dex higher than the Geneva tracks, in this paper we adopt the conservative policy of incorporating the errors on the luminosity limits in the estimations of the upper mass limits as described further below.

## 4 DISCUSSION OF RESULTS FOR SN 2001DU

### 4.1 Reddening and metallicity of the SN 2001du region

Several methods can be used to estimate the reddening towards the supernova, and hence towards the progenitor object. These have been discussed previously in Smartt et al. (2001a; 2002a) and in Leonard et al. (2002a), and here we try three different methods to determine the extinction. The method of using the Na ID interstellar absorption lines as a tracer of reddening along the line of sight is also often employed for nearby SNe. However these lines are not visible in the Sept. 9.35 spectrum. We have measured the upper limit to their strength of EW $\lesssim$ 0.4Å which is already suggestive of low interstellar absorption in the range $0.06 \lesssim E(B-V) \lesssim 0.20$ as deduced from the empirical relations between the EW of the Na ID features and line of sight reddening given in Turatto et al. (2003).

#### *4.1.1 Comparison of the continuum slope with SN 1999em and a black-body function*

As discussed in Leonard et al. (2002a) and Eastman et al. (1996), a hard upper limit to the extinction toward a SN II-P in the photospheric stage can be determined. An arbitrarily, and unphysically, hot black-body temperature is compared with the dereddened continuum of the supernova. If SN 2001du is dereddened by $E(B-V)$ >0.6 then even a function with $T_{\rm BB} = 10^9$ K is unable to reproduce the slope of the continuum (see Fig. 3). As pointed out by Leonard et al. (2002a) it is desirable to have a spectrum taken as early as possible after explosion, because as the photosphere cools the upper limit that can be derived will become less restrictive. Using this method they derived a limit of $E(B-V) < 0.45$ for SN 1999gi with a spectrum taken 0.62 days after discovery. As we discuss below, our earliest spectrum of SN 2001du was taken approximately 8 days after discovery. This establishes a very secure upper limit to the extinction, however in all likelihood the actual reddening will be lower, and a more applicable $T_{\rm BB}$ will be of the order 10000−20000 K. In Fig. 3 we plot the black-body fluxes along with suitably dereddened observed continua, and conclude that the reddening towards SN 2001du is likely to be of the order 0.0−0.22, with a hard upper limit of $E(B-V)$=0.6.



**Figure 4.** Two-colour diagram showing of the supergiants of NGC 1365, in the vicinity of SN 2001du. Also shown is the sequence of theoretical supergiant colours (Bessell 1990) reddened by $E(V-I) = 0.155 \pm 0.028$. The reddening vector is indicated by the arrow.

The supernova SN 1999em is a very well sampled nearby type II-P event and has been studied in detail by Baron et al. (2000), Hamuy et al. (2001), Leonard et al. (2002b) and Elmhamdi et al. (2003). The latter three studies have extensive optical spectroscopy during the plateau phase. Hamuy et al. have kindly made their data available in electronic format on the SUSPECT[1] archive website, and we have access to the Elmhamdi et al. electronic data. The slopes of the continua in the Elmhamdi et al. and the Hamuy et al. spectra around day +8 after explosion are very similar, and in Fig. 3 the similarity with SN 2001du is illustrated. The time of explosion of SN 1999em is taken as $t_0$ = 2451478.8 JD from the EPM analysis (Hamuy et al. 2001), hence the 1999 November 14 spectrum corresponds to day +18. Fig. 3

---

[1] http://tor.nhn.ou.edu/~suspect/

**Table 2.** Limits on the bolometric magnitude and luminosity of the progenitor star of SN 2001du using the $5\sigma$ limits from the $V_{555}$ and $I_{814}$ images.

| | | $M_V = -7.5$ | | | $M_I = -8.2$ | | |
|---|---|---|---|---|---|---|---|
| Spectral Type | $\mathrm{T_{eff}}$ (K) | BC | $\mathrm{M_{bol}}$ | $\log L/L_{\odot}$ | BC+$(V-I)_0$ | $\mathrm{M_{bol}}$ | $\log L/L_{\odot}$ |
| O9 | 32000 | −3.18 | −10.68 | 6.17 | −3.65 | −11.85 | 6.64 |
| B2 | 17600 | −1.58 | −9.08 | 5.53 | −1.78 | −9.98 | 5.89 |
| B5 | 13600 | −0.95 | −8.45 | 5.28 | −1.00 | −9.20 | 5.58 |
| B8 | 11100 | −0.66 | −8.16 | 5.16 | −0.64 | −8.84 | 5.43 |
| A0 | 9980 | −0.41 | −7.91 | 5.06 | −0.33 | −8.53 | 5.30 |
| A2 | 9380 | −0.28 | −7.78 | 5.01 | −0.14 | −8.34 | 5.23 |
| A5 | 8610 | −0.13 | −7.63 | 4.95 | 0.12 | −8.08 | 5.13 |
| F0 | 7460 | −0.01 | −7.51 | 4.90 | 0.40 | −7.80 | 5.02 |
| F2 | 7030 | 0.00 | −7.50 | 4.90 | 0.47 | −7.73 | 4.99 |
| F5 | 6370 | −0.03 | −7.53 | 4.91 | 0.55 | −7.65 | 4.96 |
| F8 | 5750 | −0.09 | −7.59 | 4.93 | 0.63 | −7.57 | 4.92 |
| G0 | 5370 | −0.15 | −7.65 | 4.96 | 0.69 | −7.51 | 4.90 |
| G2 | 5190 | −0.21 | −7.71 | 4.98 | 0.77 | −7.43 | 4.87 |
| G5 | 4930 | −0.33 | −7.83 | 5.03 | 0.78 | −7.42 | 4.86 |
| G8 | 4700 | −0.42 | −7.92 | 5.06 | 0.73 | −7.47 | 4.88 |
| K0 | 4550 | −0.50 | −8.00 | 5.10 | 0.74 | −7.48 | 4.89 |
| K2 | 4310 | −0.61 | −8.11 | 5.14 | 0.79 | −7.41 | 4.86 |
| K5 | 3990 | −1.01 | −8.51 | 5.30 | 1.09 | −7.11 | 4.74 |
| M0 | 3620 | −1.29 | −8.79 | 5.41 | 0.88 | −7.32 | 4.82 |
| M2 | 3370 | −1.62 | −9.12 | 5.54 | 0.82 | −7.38 | 4.85 |
| M5 | 2880 | −3.47 | −10.97 | 6.28 | 0.67 | −7.53 | 4.91 |

shows the excellent match between the +18 day spectrum of SN 1999em and our SN 2001du ESO spectrum taken on 2001 September 9.35. We attempted to match the SN 2001du spectra with the Hamuy et al. data for +13 (November 9) and +23 (November 19) days and applied variable reddening to the spectra of SN 2001du but could not consistently match the continuum slope, the strength of the absorption lines and the H$\alpha$ P-Cygni feature. In addition we varied the absolute linear scaling factor applied to the SN 2001du spectrum (to account for distance and intrinsic magnitude differences) and the amount of dereddening (assuming R=3.1) when matching it to the +18 day SN 1999em spectrum. We found that any extra reddening added to SN 2001du was incompatible with a good match and even applying an additional reddening of $E(B-V) = 0.1$ gave poor results. Hence the spectrum of SN 2001du taken on 2001 September 9.35 UT is virtually identical with that of SN 1999em at +18 days after explosion and implies an identical reddening. Baron et al. (2000) have compared the early time optical and UV spectra of SN 1999em with a non-LTE model atmosphere fit. They simultaneously determine a temperature and a low value of reddening of $E(B-V) = 0.10 \pm 0.05$, and argue that increasing the value of $E(B-V)$ will not lead to a consistent fit to all the features in the spectra. In summary, the continua and line strengths of SN 1999em and SN 2001du on +18 days are virtually identical, and any further artificial reddening of SN 2001du leads to significant inconsistencies. Hence our estimate of reddening towards SN 2001du using this method is $E(B-V) = 0.1 \pm 0.05$.

### *4.1.2 Estimates of reddening from nearby H* II *regions*

The abundance gradient in NGC1365 has been studied by Roy & Walsh (1997) who observed the nebular lines of 55 H II regions. For each H II region they determined an extinction at H$\beta$, which is the combined value of Galactic reddening ($A_V = 0.16$), and extragalactic extinction. Their nebular region RW21 is virtually coincident with the star formation region around the position of the supernova, and the centre of this H II region is estimated to be approximately 4″ from SN 2001du. Hence it is likely that this nebulosity is directly associated with the ionizing hot stars in the vicinity of SN 2001du i.e. the bright clustered sources in the $U_{336}$ image in Fig. 1(a). Roy & Walsh (1997) measured a logarithmic extinction at 4861Å of $c(\mathrm{H}\beta) = 0.83 \pm 0.06$ for RW21 and using an extinction law of $A(4861)/A(V) = 1.16$ (Cardelli et al. 1989) this translates into $A_V = 1.8 \pm 0.1$, or $E(B-V) = 0.58 \pm 0.03$ (assuming R=3.1). There are two other H II regions lying in the same spiral arm at the end of the central bar in NGC1365 both at distances of approximately 15″ from the supernova (RW19 and RW20). These have a virtually identical extinction of $E(B-V) = 0.53 \pm 0.03$, hence the extinction toward the ionizing regions near the SN position which produce the nebular lines is of order $E(B-V) = 0.58 \pm 0.03$. However as discssed by Calzetti et al. (1994) and Calzetti (1997) the extinction measured in starforming regions from the nebular lines is generally a factor of two higher than that of the underlying stellar continuum. Hence the H II region reddening is not likely to be directly applicable to the SN 2001du reddening as the supernova it is not coincident with the position of the UV-bright ionizing O-stars. We hence assume that a reddening a factor of two lower should be applicable to the stars near the

RW21 region i.e. $E(B-V)=0.29\pm0.03$. While this may appear somewhat arbitrary, we should get a reliable mean when combining it with the other extinction estimates .

Roy & Walsh determined the oxygen abundance in their H II regions from an analytic calibration of the $R_{23}$ ratio $R_{23}$=([OII]$\lambda$3727+[OIII]$\lambda\lambda$4959,5007)/H$\beta$. While this method usually leads to a reasonably consistent estimation of how abundances differentially vary within a galaxy, the absolute abundance determined for any particular $R_{23}$ ratio is highly dependent on the calibration employed. Recently Smartt et al. (2001b) and Trundle et al. (2002) have discussed various literature calibrations of the $R_{23}$ ratio, and compared extragalactic nebular abundances directly with massive stars born within the H II regions. They suggest that the calibrations of Pilyugin (2002) and McGaugh (1991) (the latter parameterized by Kobulnicky et al. (1999)) give the best agreement for oxygen abundances in the gas and in massive B-type supergiants. Applying these calibrations to the $R_{23}$ ratio of RW21 results in an oxygen abundance $12+\log$ (O/H)=8.5 dex (from the Pilyugin calibration), and 8.8 dex (from McGaugh). Hence within the uncertainty of the absolute calibration of this method, the H II region RW21 is approximately solar in metallicity; assuming the solar oxygen abundance is 8.83 dex (Grevesse & Sauval 1998).

### *4.1.3 The reddening towards surrounding stars in NGC1365*

In this case the reddening can be measured from 3-colour photometry of the bright supergiants in the area around the supernova, and one then assumes that this is representative of the reddening toward the supernova itself. Photometry in three filters (F555, F675W and F814W transformed to standard $VRI$) was measured for 190 stars on the PC chip in the vicinity of SN 2001du (to a radius of 23″from the SN position). The locus of the observed stars, in a two-colour plane $(V-I, V-R)$ was compared with a theoretical supergiant colour sequence from Bessell (1990) for galactic supergiants. The reddening was determined by ascertaining the displacement between the observed stars and the theoretical sequence in the two-colour plane. A weighting factor was included in the calculation, in favour of those stars with more accurately determined magnitudes. The reddening was determined to be $E(V-I)=0.16\pm0.03$. Fig. 4 shows the two-colour diagram of the stars imaged in the vicinity of SN 2001du; also shown is the theoretical supergiant sequence appropriately reddened. Utilising a reddening law of $R_V=A_V/E(V-I)=2.45$ (Silbermann et al. 1999) yielded a value of $A_V=0.38$. In determining a reddening towards the Cepheid population of NGC1365 (in order for distance determinations), Silbermann et al. (1999) measured a value of $A_V=0.40$. Within the errors this is identical to our measurement in the field of SN 2001du, and suggests that the reddening towards this part of the galaxy is not significantly different to that derived by Silbermann et al., and is not prohibitively high. A standard reddening law of $R_V=A_V/E(B-V)=3.1$ would imply a value of $E(B-V)=0.12\pm0.02$ is appropriate for the line-of-sight to SN 2001du from this method.

## 4.2 A luminosity and mass limit for the progenitor of SN 2001du

The three methods above give reassuringly consistent measurements for the reddening towards SN 2001du, and presumably its progenitor star. Hence we adopt a mean value of these estimates of $E(B-V)=0.17\pm0.09$ and (assuming $R_V=3.1$) $A_V=0.53\pm0.26$. Smartt et al. (2001a; 2002a) discuss the general technique employed to determine an upper luminosity limit for the progenitor for it not to have been detected on pre-explosion images. The pre-explosion $U_{336}$ archive image was not sufficiently deep to place any significantly constraining upper luminosity limits for the progenitor star and is not considered further in the following discussion. The $V_{555}$ and $I_{814}$ $5\sigma$ detection limit magnitudes were transformed to $V=24.4$ and $I=23.5$ using the transformations of Holtzman et al. (1995b). The corresponding absolute magnitudes ($M_V$ and $M_I$) were calculated by correcting for the extinction, determined earlier, and adopting a distance modulus to NGC 1365 of $(m-M)=31.39\pm0.26$ (Ferrarese et al. 2000). The limiting absolute magnitudes were determined to be: $M_V=-7.5$ and $M_I=-8.2$. Bolometric magnitudes, for each waveband, were determined by applying bolometric corrections, and an appropriate colour correction $(V-I)_0$ for $M_I$, both taken from Drilling & Landolt (2000). The luminosity limits, as a function of temperature, were calculated from the bolometric magnitudes assuming a solar bolometric magnitude of $M_{\rm bol}=+4.74$, and are listed in Table 2. These luminosity limits were overlaid on an HR diagram with our evolutionary tracks (as described in Section 3). An HR diagram showing the exclusion region defined by the luminosity upper limits is shown in Fig. 5. A closer view of the end-points of the tracks in the red supergiant region, and the exclusion limit defined by the $I$ band sensitivity limit is further shown in Fig. 6. The expected magnitude errors in distance modulus (±0.26), reddening (±0.26), sensitivity limits (±0.2), and BC and intrinsic stellar colours (±0.4) are combined in quadrature giving an uncertainty on $M_{\rm bol}$ of ±0.58 mag. As we pointed out in Section 3, the end-points of our tracks in the red-supergiant region tend to be systematically higher than the Geneva tracks by 0.1−0.15 dex. As the observational uncertainties are significant we adopt the conservative approach of adding the total error to the luminosity upper limit to derive a robust upper mass-limit. In this case we can say that the upper limit to the initial mass of the progenitor star of SN 2001du is 15M$_\odot$.

As discussed in Sect. 2.1 we believe it unlikely that star B is a single point source which is precisely coincident with the the supernova and is an unlikely progenitor candidate. However if it was the progenitor it would have $\rm M_V=-6.8$, and $V-I\leq1.4$. This would constrain it to be just below the exclusion region in Fig. 5 and to be bluer than approximately K2 spectral type i.e. cooler than $\log T_{\rm eff}\lesssim3.7$, and with a mass in the region 15-20M$_\odot$. Hence it would not have been an M-type supergiant, but an unperturbed early K-type could provide a large enough hydrogen envelope to sustain a plateau phase of ∼100 days (see Table 3), and is thus not inconsistent with the supernova evolution. Van Dyk et al. (2003b) have also reanalysed the site of SN 2001du using our images of the supernova, and come to the same conclusion that Star B is not the progenitor. The upper mass



**Figure 5.** SN 2001du: Luminosity limits from pre-explosion images in the $V$ and $I$ are plotted as thick solid lines. The shaded region is where a progenitor would have been detected in at least one filter. Overlaid are the evolutionary tracks as described in Sect. 3 for stars with main sequence masses $9-60\mathrm{M}_{\odot}$, for solar metallicity. The locations of the progenitors of SN 1987A and SN 1993J are also shown.



**Figure 6.** SN 2001du: End stages of the evolutionary tracks for stars with initial masses in the range $9-20\mathrm{M}_{\odot}$. The limiting luminosities are shown as the thick solid lines. Uncertainty limits, on the I-band limiting luminosity are shown as the dashed lines. The limiting luminosity including the expected error constrains a robust upper mass limit of $15\mathrm{M}_{\odot}$



**Figure 8.** SN 1999em: updated exclusion region diagram for the progenitor of SN 1999em (from combined $R$ and $I$ limits), using the new distance estimate for NGC1637 of 11 Mpc. Our evolutionary tracks as discussed in Sect. 3 are plotted along with the uncertainties on the luminosity limits. The new distance increases the upper mass limit to $15\mathrm{M}_{\odot}$.

limits derived, again repeating our method, are in agreement with those derived here $(13\mathrm{M}\gtrsim^{+7}_{-4})$, although we have reduced the uncertainties using our finer grid of stellar evolutionary tracks.

## 5 DISCUSSION OF RESULTS FOR TWO OTHER II-P SUPERNOVAE

### 5.1 SN 1999em

In Smartt et al. (2002a) we discussed the prediscovery images of SN 1999em which were taken from the ground with the Canada-France-Hawaii telescope at a resolution of $0.7''$. As with SN 2001du and SN 1999gi (see Section 5.2) there was no detection of a progenitor star at the position of SN 1999em, and the sensitivity limits of the $VRI$ frames allowed an upper mass limit to be derived. Similar methods as discussed above led us to determine an upper mass limit of approximately $12\mathrm{M}_{\odot}$. However these conclusions are based on the assumption that the ground-based images do not have contaminating objects within the seeing PSF of $0.7''$. As shown in the case of SN 1999gi, which exploded in a fairly large OB-association, it is quite possible that ground-based resolution images would not be adequate to resolve the individual massive stars. For that reason we have reobserved the environment of SN 1999em with WFPC2 during our Cycle 10 program GO9041. The main goal of this observation was to investigate the immediate environment surrounding the progenitor of SN 1999em, and determine if a contaminating star cluster would compromise the result. SN 1999em was observed with WFPC2 on 2002 January 15 through the filters F555W, F675W and F814W. The F555W image is shown in Fig. 7, and we have identified the position of the supernova through matching the brightest single stars in the field of the WFPC2 data and the WHT image of Smartt et al. (2002a). A geometric transformation between the two frames results in a residual uncertainty of approximately $0.1''$ in the SN position. The WFPC2 image shows a faint object at the position of the supernova, which is not a clean point source. Two peaks appear, superimposed on a quite faint extended region of flux. Although this diffuse emission is faint it is clearly visible in both the F555W, and the F675W but not the F814W filter. We estimate the position of the brighter peak at the centre of Fig 7a is within $0.08''$ of the supernova centroid, which is within the errors of our differential astrometry. The extended weak flux is approximately $0.5''$ in diameter, or 25 pc (assuming the distance discussed below). This is much too far to be due to a light echo from SN 1999em and is likely a small star cluster which hosted the progenitor star. The magnitude of this object as a whole (from aperture photometry using a radius of $0.3''$) is $V_{555} = 23.7 \pm 0.2$, which is a combination of the residual supernova magnitude and the underlying association. This would imply that the host cluster



**Figure 7.** **(a):** A WFPC2 F555W image of the SN 1999em taken on 2002 January 15, some +810 days after explosion. The position of the supernova was identified from the high-resolution WHT images presented by Smartt et al. (2002a). **(b):** Close up of the faint source at the position of SN 1999em, which is not a simple point source (compare with the other two point sources at $y \simeq 0.6$). It appears as two peaks superimposed on a faint diffuse background. However the region around the SN position is uncrowded and relatively clear of contaminating bright sources within the $0.7''$ seeing disk of the pre-explosion frames of Smartt et al. (2002a). This validates the use of these lower resolution frames in setting a luminosity and mass limit for the progenitor star.

has $M_V > -6.1$, which is below the detection limit of the pre-explosion frames ($M_V = -6.49$; Smartt et al. 2002a). Hence the size and luminosity of the cluster is fairly typical for the smallest young associations found in spirals and in the Local Group e.g. Bresolin et al. (1996; 1998). For this paper, the most important conclusion is that the images confirm that within the seeing disk of our prediscovery CFHT images there are no other *bright* contaminating objects.

Leonard et al. (2003, in preparation) have observed the galaxy NGC1637 on several epochs with HST/WFPC2 during Cycle 10 to measure a Cepheid distance to the galaxy. This is the first attempt to compare a Cepheid distance to the expanding photosphere method as applied to the type II supernovae, and the Cepheid distance measurement is $D \sim 11$ Mpc (D. Leonard and N. Tanvir, private communication). Hence we have revised the exclusion diagram presented in Smartt et al. (2002a) with this new distance. As the HST/WFPC2 images confirm that no contaminating stellar sources exist within the ground-based seeing disk that would compromise the point source sensitivity limits, the diagram shown in Fig. 8 should now be a robust result. Applying the error estimates and (as in Smartt et al. 2002a) the same conservative criteria as applied to SN 2001du of using the highest mass that falls within the upper bounds of the errors, we now derive an upper mass limit of $15\mathrm{M}_{\odot}$. The fact that the underlying association or cluster, in which the progenitor was born, was small adds further weight to our argument that the progenitor could not have been very massive. Assuming a typical initial mass function, it is statistically unlikely that a cluster of this size would have given birth to very massive stars of greater than $20\mathrm{M}_{\odot}$. The new limit derived is tabulated in Table 3 for comparison with other known progenitors.



**Figure 9.** SN 1999gi: End points of stars with initial masses in the range $8{-}20\mathrm{M}_{\odot}$ using our evolutionary tracks as described in Sect. 3. The limiting luminosities (from the F606W filter) are shown as the thick solid lines and the uncertainties on these are the dashed lines. This gives a robust upper limit of $12\mathrm{M}_{\odot}$.

The unexpected faint magnitude of the supernova has implications for the Elmhamdi et al. (2003) conclusions. Even assuming the total magnitude measured (i.e. supernovae plus underlying cluster) to be an upper limit to the SN 1999em flux itself, it is significantly fainter than the last reported detection by Elmhamdi et al. (2003) of $V = 20.05 \pm 0.18$ on 2001 March 16. Elmhamdi et al. have suggested that their last photometric point is fainter than the expected linear decay (extrapolated from previous points) of $\gamma_v \sim 0.97$ mag $(100\mathrm{d})^{-1}$ by approximately 30%,



**Figure 10.** Estimate of the lower mass limit for the progenitor of SN 1997bs. The object was detected at $M_V \simeq -8.1$ (in F606W; Van Dyk et al. 2000), and the bolometric luminosity locus as a function of $T_{\rm eff}$ (from the supergiant table in Drilling and Landolt 2000) is plotted. Overlaid are evolutionary tracks for stars with main sequence masses $12-40{\rm M}_{\odot}$, for solar metallicity. The progenitor must have been somewhere on this line prior to explosion, hence an approximate lower limit to the mass is M> $20{\rm M}_{\odot}$.

and have interpreted this as a sign of dust formation. If we extrapolate the linear decay of Elmhamdi et al. (from the last point on 2001 February 01), then the SN 1999em should be V≃23.0 (at day +810) in the HST WFPC2 images. As it is $0.7^m$ fainter (assuming the F555W bandpass is close to $V$) this is supporting evidence that the deviation they reported is real, and continues even more severely at late times. Elmhamdi et al. also suggested that the [O I] line profile variations after day ∼465, were indicative of dust formation in the cooling supernova. This faint WFPC2 magnitude of the SN 1999em would seem to add further support to that. The multi-epoch images of Leonard et al. (G0 9155) to monitor the Cepheid variable population also contain the SN 1999em on the field of view, hence this will allow them to probe the late time lightcurve in F555W and F814W, and dust formation scenario, in much more detail.

### 5.2 SN 1999gi

In Smartt et al. (2001a) we presented the HST archive pre-explosion images for SN 1999gi along with images of the supernova taken 14 months later in order to astrometrically determine the supernova position precisely. There was no detection of a progenitor star in either the F300W or F606W filter, and using the methods described above in Section 4.2 we derived an upper mass limit for the progenitor star of $9^{+3}_{-2}{\rm M}_{\odot}$, which assumed it was a red supergiant in the final stages of its evolution. However in this paper we assumed a distance of $7.9\pm2$ Mpc, and recently Leonard et al. (2002a) have determined an improved distance to SN 1999gi using the expanding photosphere method. They estimate a distance of $11.1\pm2$ Mpc, and have also carried out a careful comparison of reddening towards the supernova from four different methods (which we followed in Sect. 4.1). Leonard et al. estimate a slightly higher value of $E(B-V)=0.21$ compared to the $E(B-V)=0.15$ used in Smartt et al. (2001a), and derive an upper mass limit for the star of $15^{+5}_{-3}{\rm M}_{\odot}$ on this basis.

We have recalculated the luminosity and mass limits using the new distance of Leonard et al., and the higher reddening, but using the better sampled tracks at our disposal we can reduce the uncertainty in the mass range of Leonard et al.'s result. Leonard et al. have estimated the mass limit by adding the upper error uncertainty and simply taking the nearest modeled progenitor mass to that, with an upper limit of the next highest mass model in the Geneva tracks. The exclusion region of the HRD is plotted in Fig. 9, and applying the same criteria as for SNe 2001du and 1999em we get an upper mass limit of $12{\rm M}_{\odot}$, which is below our best estimate *and* the uncertainty. The difference between this limit and the $15^{+5}_{-3}{\rm M}_{\odot}$ estimate by Leonard et al. is that our tracks end at slightly higher luminosities than the Geneva models, and also we can rule out the rather large range by employing the integer sampled mass tracks. It is understandable that Leonard et al. employed a conservative approach in estimating the upper limit and errors. However we can certainly rule out the progenitor being a $20{\rm M}_{\odot}$ red supergiant, as this lies significantly above the detection limits – it would have a luminosity of $\log L/L_{\odot}\simeq5.3$, and hence a magnitude of $V_{606}\simeq23.9$, which would have been a clear ($10\sigma$) detection on the pre-explosion images. In addition there may be a slight discrepancy between our temperatures of pre-supernova M-supergiants and those in Fig. 10 of Leonard et al., where the effective temperatures of the evolutionary track endpoints appear too cool by ∼0.05 dex. They use the Geneva tracks (as employed in Smartt et al. 2001a), but the plot does not exactly match the tabulated values i.e. their end-points would be close to $\log {\rm T}_{\rm eff}$=3.5 in our Fig 9. If the tracks are shifted to the tabulated temperature then their luminosity limit of the F606W filter would no longer skim the $20{\rm M}_{\odot}$ track, and the $15{\rm M}_{\odot}$ track would be inside the exclusion region.

We accept the distance and reddening determinations of Leonard et al. as more appropriate to those used in Smartt et al. (2001a), and that Leonard et al. are being understandably cautious in using the estimate plus uncertainty to derive a very hard upper limit. But the simplest interpretation of Fig. 9, where we use better sampled end points in the tracks, indicates an upper mass-limit of $12{\rm M}_{\odot}$. Clearly these assumptions rely on how accurate the theoretical stellar evolutionary models are in predicting the luminosity and temperatures of M-type supergiants.

### 5.3 Progenitors of core-collapse supernovae

In this section we review all of the direct observational evidence that exists for core-collapse precursors. The events for which we have some restrictive information on the progenitors are listed in Table 3. Only those supernovae which are fairly certain to be core-collapse explosions, *and* have a well defined SNe type attributed are included.

The event SN 1978K is not included although it had a progenitor object visible. The supernova itself was serendipitously discovered in January 1990 as a peculiar object on an archive photographic plate (Ryder et al. 1993). Hence there is only a crude light-curve from archive plates taken covering that region of sky. Ryder et al. (1993) conclude that it was more likely to be a type II event rather than an LBV

**Table 3.** A list of known supernova progenitors and constraints on their masses and spectral types for supernovae which are certain to be real core-collapse events and have been reliably typed. The estimated initial mass of the progenitor star is listed as $M_i$ and is from direct measurements of the progenitor star or limits thereon. Compilation values for mass and metallicity are taken from Smartt et al. (2002a), and updated with results from this paper and Smartt et al. (2002b; for SN 2002ap). We estimated the metallicity of the progenitor of 1997bs simply from its distance from the centre of NGC3627, and applied a typical abundance gradient for an Sb galaxy from Vila-Costas & Edmunds (1992). The mass of the ejecta in the supernova is $M_{ej}$ and the radii of the progenitors are estimated from analysis of the supernova spectra and lightcurve, as described in Section 5.3. The values for SN 1987A and SN 1993J, and SN 1980K are taken from Arnett (1996) and Woosley et al. (1994).

| SN | Galaxy | Distance (Mpc) | Type | Metallicity ($Z/Z_\odot$) | Progenitor properties<br>Type | <br>$M_i/M_\odot$ | <br>$M_{ej}/M_\odot$ | <br>$R/R_\odot$ |
|---|---|---|---|---|---|---|---|---|
| 2002ap | NGC628 | 7.3 | Ic | 0.5 | WR? | <40 | $2.5-5$ | – |
| 1997bs | NGC3627 | 11.4 | IIn | ∼1 | ? | >20 | – | – |
| 1987A | LMC | 0.05 | II-pec | 0.5 | B3Ia | 20 | 15 | 43 |
| 1993J | M81 | 3.6 | IIb | 2 | G5−K0 Ia | 17 | 3 | ∼500 |
| 1980K | NGC6946 | 5.1 | II-L | 0.5 | ? | <20 | 2 | ∼300 |
| 2001du | NGC1365 | 17.9 | II-P | ∼1 | G-M type | <15 | – | – |
| 1999em | NGC1637 | 11 | II-P | 1-2 | K-M type | <15 | 5−18 | 260−1500 |
| 1999gi | NGC3184 | 11.1 | II-P | ∼2 | G-M type | <12 | 10−30 | 100−400 |



**Figure 11.** The theoretical SNe populations from the models of Heger et al. (2003), but ignoring their extremely metal poor and metal-free objects. All objects below the dotted line should be weak SN due to black-hole formation by fall-back of most of the material surrounding the iron core. All objects to the right of the solid black line should have lost their hydrogen envelope at the point of core-collapse, and hence should be hydrogen free SNe. The blank white regions are where no supernova is seen; below $9M_\odot$ due to the formation of white dwarfs in low mass stars; above $40M_\odot$ due to direct black hole formation. The observed positions of the supernovae progenitors in Table 3 are approximately estimated and placed, with arrows indicating those with limits. To place the stars in the metallicity dimension we have assumed a zero-point on the y-axis of $\log Z/Z_\odot$=-1.

"super-outburst", however this is not beyond doubt. If it was a supernova we do not know its type, peak magnitude or very much about the lightcurve. Given the limited resolution of the photographic surveys (particularly close to the plate limit) we do not know if the object at the position of SN 1978K is really a point source, a nebula or cluster (Ryder et al. quote a distance of 4.5 Mpc). These uncertainties lead us to omit SN 1978K from Table 3 for now, at least until some of the questions can be definitively answered. The event SN 1961V is also not included for similar reasons. It is very unclear if it was a genuine supernova or an S-Doradus outburst from a very massive star like $\eta$ Carinae (Humphreys et al. 1999). The progenitor object was possibly identified as a star with an absolute magnitude $M_{\rm bol} \simeq -11$, which is much too bright for a stable massive star or $\eta$ Car like object in a quiescent state. Hence it was likely to be in outburst during its period as an $M_{\rm bol} \simeq -11$ object, and given the sparse data available we do not have a definitive model for the physics behind the eruption. Again because of these uncertainties, we do not discuss this further.

A detection of a progenitor star associated with SN 1997bs in M66 has been claimed by Van Dyk et al. (2000), who originally suggested that this event was also not a genuine core-collapse supernovae but another super-outburst similar to SN 1961V and $\eta$ Car, and that the progenitor actually survived the explosion in HST images. However Li et al. (2002) have observed SN 1997bs at a later epoch with HST and observed it to fade considerably, and get bluer, compared to the Van Dyk et al. magnitudes. With this new information, the conclusion is that SN 1997bs was indeed likely to have been a subluminous IIn supernova, and the progenitor has not survived. Van Dyk et al. (2000) have estimated the absolute magnitude of the progenitor star to be $M_V \simeq -8.1$ from the single exposure F606W WFPC2 image. As no colour information is available we do not know the temperature or appropriate bolometric correction to allow the progenitor to be approximately placed on an HR diagram. However given this detection in a single band, a *lower* limit can be placed on the mass of the progenitor. In Fig 10 we plot the locus of a $M_V = -8.1$ star as a function of effective temperature and hence bolometric correction. The progenitor must have existed at some point on this locus, hence allowing a lower limit to its mass of approximately $20{\rm M}_{\odot}$.

The brightest SNe Ic to occur in recent times was SN 2002ap in M74 (=NGC628), and its early lightcurve and spectral evolution was recently studied by Mazzali et al. (2002). By comparing explosion models of CO cores combined with spectral synthesis to model the optical observational data, they estimated an ejecta mass of between $2.5{-}5{\rm M}_{\odot}$. With an assumed (but uncertain within a factor $\sim$2) remnant mass of roughly $2.5{\rm M}_{\odot}$, this would imply the mass of the CO precursor star should be in the conservative range $2.5{-}10{\rm M}_{\odot}$. The final masses of Wolf-Rayet stars of the WC type in the LMC have been estimated by Crowther et al. (2002) to be in the range $11{-}19{\rm M}_{\odot}$, and those in the galaxy in the range $7{-}14{\rm M}_{\odot}$. Given the uncertainties, there is reasonable agreement between these two estimates suggesting a WC type progenitor (of initial mass $\sim30{\rm M}_{\odot}$) is consistent with the characteristics of SN 2002ap. Deep ground-based images of the pre-explosion site of the supernova were presented by Smartt et al. (2002b), and no detection of a massive star was found. This paper ruled out most of the upper HR-diagram as possible progenitor sites, although this is to be expected as SNe Ic show no evidence of the progenitors having H-rich atmospheres. It is likely that the precursor was a WR star in the WC phase, and had an initial mass of less than $\sim40{\rm M}_{\odot}$. However initially lower mass stars which have had their H-envelopes stripped during mass-transfer in a binary system, rather than through a radiatively driven stellar wind, still do remain viable precursors.

The three II-P supernovae discussed above (SNe 2001du, 1999em, 1999gi) are also listed in Table 3, along with a similarly determined upper mass limit for SN 1980K. Pre-explosion images for this supernova were presented by Thompson (1982), and we have recalculated the limit using the same evolutionary tracks as employed above. The more definite values for SN 1987A and 1993J are included. The analytical model for the plateau stage of type II supernovae of Popov (1993) produces equations relating the observed quantities of $t_{\rm p}$ (the plateau duration), $M_V$ (absolute visual magnitude at a representative point during the plateau), and $u_{\rm phot}$ (the expansion velocity of the photosphere) to three intrinsic properties of the supernova. These are energy of the explosion and the mass and radius of the initial envelope of the progenitor star. More detailed hydrodynamic models of SNe II-P by Litvinova & Nadyozhin (1985) have produced similar relations, but with different coefficients. The mass and initial radius of the ejected stellar envelope for 1999em and 1999gi has been estimated by Elmhamdi et al (2003), Hamuy (2003) and Nadyozhin (2003) using both sets of equations but differing values for $t_{\rm p}$, $M_V$ and $u_{\rm phot}$. The mass and radii derived in each paper differ quite substantially, depending on what values are assumed. The main differences are the distance assumption, which critically affects $M_V$, and the epoch at which $u_{\rm phot}$ is measured. We have recalculated the range in feasible values using the most up to date values for distance and reddening (as discussed above), and get values of $M_V = -16.7$ (1999em) $M_V = -15.9$ (1999gi) at approximately 50 days after discovery (i.e. in the middle of the plateau). We assume that $t_{\rm p} \simeq 100$ days and vary $u_{\rm phot}$ between $3000{-}4000$km s$^{-1}$ . There appears no discrepancy in the actual measurements of $u_{\rm phot}$ in 1999em between Elmhamdi et al. (2003), Hamuy et al. (2001) and Leonard et al. (2002b), but there is some difference as to which epoch is chosen to measure $u_{\rm phot}$ to use in the calculations. The ranges of ejected mass (${\rm M}_{ej}$) and progenitor radius (R) are tabulated in Table 3. If we allow $\sim2{\rm M}_{\odot}$ to be lost by mass-loss and neutron star formation, then the mass ranges for the envelope are in reasonable agreement with our direct estimates for the initial mass of the progenitor stars. The large range in radius of the envelope is consistent with spectral types between G0−M0. We do not have enough photometric information available for SN 2001du to estimate these values for comparison. However our direct mass limit for SN 2001du is similar to those for SNe 1999em and 1999gi. Given the very similar spectra of SN 2001du with both of the latter (see Fig. 3), this is not altogether surprising. This supports our previous suggestion (Smartt et al. 2002a) that normal SNe II-P result from fairly low mass stars ($<12{\rm M}_{\odot}$) with large radii in the M-supergiant phase that have undergone only moderate mass-loss. The fact that 1999em and 1999gi were faint in the x-ray and radio suggests that their progenitor stars had undergone fairly moderate or low mass-loss. The fluxes in these wavebands have allowed Schlegel (2001) and Pooley et al. (2002) to estimate mass-loss rates for the progenitors of these two supernova which are the order of $\dot{M} \sim 10^{-6}\mathrm{M}_{\odot}\,\mathrm{yr}^{-1}$, with a conservative uncertainty of a factor of two. This supports our result that the progenitors were not very massive stars, as one would expect such stars to have gone through more drastic mass-loss events and hence to have produced higher x-ray and radio fluxes. Hence we conclude that moderate mass stars (of less than approximately $15\mathrm{M}_{\odot}$) which end their lives in the red supergiant phase tend to give rise to a fairly homogeneous class of SNe II-P , and initially higher mass stars likely give rise to more heterogeneous supernovae, although clearly more results are needed to increase the statistics.

Recent theoretical work by Heger et al. (2003a; 2003b) on the the pre-supernova evolution of massive stars as a function of mass and metallicity, has produced supernova population diagrams. These relate the initial mass of a star to the type of supernova that is produced, and how that roughly depends on metallicity of the original star. The metallicity plays a key role in determining the mass-loss rate, which in turn is a critical factor in the evolution. This has initially been done for single stars, with no rotation considered. We have reproduced the figure of Heger et al. (2003b), and ignored the very metal poor and metal-free regions as they are not applicable to any of the local Universe supernovae discussed above. The regions between the II-P, II-L and Ib/c regions are delineated by the mass of the hydrogen envelope which is left just before core-collapse, which in turn is dependent on mass-loss history and hence metallicity. In Fig. 11 we have put the positions of the supernovae listed in Table 3 to compare with the model predictions. We emphasise that at present this is a rather qualitative comparison given the approximate metallicity scale proposed on the ordinate, and the limits we have for the various nearby supernovae. Nevertheless it is certainly valid and essential to compare the state-of-the-art observational and theoretical results in this data starved area. The positions of the three SNe II-P are in good agreement with the models, as is the position of SN 1987A. However SN 1980K was a type II-L but appears in the theoretical II-P region. Even if the metallicity is uncertain by a factor of two, it would be difficult to envisage pushing this into the II-L region as it stands. This suggests that the progenitor lost more mass than one would expect for a $20\mathrm{M}_{\odot}$ star of metallicity between $0.5{-}1\mathrm{Z}_{\odot}$. The extra mass-loss could be accounted for by interaction within a binary system, with the progenitor losing its mass through mass-transfer rather than radiatively driven winds. SN 1997bs is also likely to have been too massive for the II-P region, although given that we have only a *lower* limit, the actual mass could sit comfortably in the II-L area. One serious discrepancy is obvious in that SN 2002ap was a bright Ic supernova, however the models suggest that these only come from either metal-rich stars, or very massive stars with $M_i > 60\mathrm{M}_{\odot}$. As discussed in Smartt et al. (2002b) the initial mass is unlikely to have been as high as $\sim 60\mathrm{M}_{\odot}$, hence there is a discrepancy with the position of the weak SN region. Smartt et al. (2002b) estimated the metallicity to be roughly $0.5\mathrm{Z}_{\odot}$, and again even if this was solar the position of SN 2002ap would still be discrepant with the normal SN Ib/c population region. From the Heger et al. figure one would expect that bright Type Ib/c only come from massive single stars if the metallicity is quite high, and hence they should be relatively rare. However invoking mass-loss again through close binary evolution at low metallicities could account for objects such as SN 2002ap. The future statistics of masses and mass limits for the progenitors of SNe Ib/c will be of crucial importance in distinguishing between the two scenarios. In summary then it would appear that the theoretical II-P region stretches to masses which are a little too high, and that the weak SN region should be confined to lower metallicities or much higher masses. This is of course a very preliminary comparison, with the caveat that only single stellar evolution is included in the supernova population diagram. The prospects of setting more constraining limits on SNe progenitors or detecting the stars themselves will allow better observational constraints to be placed on these models in the future (see Smartt et al. 2002a for a discussion of future prospects).

## 6 CONCLUSIONS

We have investigated HST images of the site of the nearby supernova SN 2001du taken 6.6 years before explosion, and compared our results with similar pre-explosion investigations of other core-collapse supernovae. Our findings are summarised below

(i) Although there is a $3\sigma$ detection of a source very close to the supernova position in the WFPC2 $V-$band image, it is not precisely coincident with the supernova centroid. It is slightly beyond the error radius in our analysis and is not a clear detection of an unambiguous point source. We conclude that the progenitor star was likely below the detection limit of the pre-discovery images and is not visible on any of the frames. Using the $5\sigma$ sensitivity limits of the images we estimate that the progenitor likely had an initial mass of $<15\mathrm{M}_{\odot}$.

(ii) SN 2001du was a type II-P, and probably very similar to two other nearby recent SNe II-P 1999gi and 1999em. We have previously set similar luminosity and mass limits on the progenitors of those events (Smartt et al. 2001a; Smartt et al. 2002a) and in this paper we have revisited those using new data. A new distance to NGC3184 determined by Leonard et al. (2002a) slightly changes the mass limit for SN 1999gi to approximately $12\mathrm{M}_{\odot}$. New HST images of the site of SN 1999em confirm the validity of our previous results, and using another new distance increases this upper mass to approximately $15\mathrm{M}_{\odot}$.

(iii) Hence all three SNe had very similar mass progenitor stars, which is not a surprising result given the photometric and spectroscopic similarity of the supernovae themselves. The results are consistent (but not uniquely definitive) with the idea of SNe II-P arising in red supergiant progenitors of moderate mass.

(iv) We have made first attempts at comparing all the known, direct, information available on supernova progenitors with theoretical models of stellar evolution up to the core-collapse stage. These models have produced supernova population diagrams which relate the initial mass and metallicity of massive stars to the types of supernovae that they produce. We find encouraging reasonable agreement between the observational and theoretical results for the masses of

II-P supernovae progenitors. However we find discrepancies for a II-L and bright Ic event. This is preliminary work that should increase in statistical significance over the coming years

## ACKNOWLEDGMENTS

This paper was based in part on observations made with the NASA/ESA *Hubble Space Telescope*, obtained from the data archive of the Space Telescope Institute which is operated by the Association of the Universities for Research in Astronomy Inc., under NASA contract NAS5-26555. These observations are associated with proposal GO9041. Some observations were collected at the European Southern Observatory, Chile, ESO67.D-0594. We acknowledge the support given by ASTROVIRTEL, a Project funded by the European Commission under FP5 Contract No. HPRI-CT-1999-00081. We thank the ASTROVIRTEL team at ESO/ST-ECF for the new software developed. SJS and JRM thank PPARC for financial support in the form of an Advanced Fellowship award and studentship respectively and CAT thanks Churchill College for a fellowship, SB acknowledges support from the Italian Ministry for Education, University and Research (MIUR) through grant Cofin MM2001021149-002. We thank Peter Meikle for useful discussions and help in the initial typing of SN 2001du, Bill Januszewski for help in coordinating and executing our HST observations, and Doug Leonard and Nial Tanvir for communicating their new Cepheid distance determination to NGC1637 prior to submission of their paper. Use was made of VSNET data for the optical light-curve of SN 2001du, for which we are very grateful.

## REFERENCES

Aldering G., Humphreys R.M., Richmond M., 1994, AJ, 107, 662
Alexander D. R., Ferguson J. W., 1994, ApJ, 437, 879
Anders E., Grevesse, N., 1989, Geochim. Cosmochim. Acta, 53, 197
Arnett D., 1996, Supernovae and Nucleosynthesis, Princeton University Press
Baron E., et al, 2000, ApJ 545, 444
Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1995, ApJ, 446, L35
Bessell M.S., 1990, PASP 102, 1181
Branch D., 2003, Proc of IAU Symp. No. 212 (2002), eds. K. A. van der Hucht, A. Herrero, C. Esteban
Bresolin F., Kennicutt R.C. Jr., Stetson P.B., 1996, AJ, 112, 1009
Bresolin F., et al. 1998, AJ, 116, 119
Branch D., Livio M., Yungelson L. R.; Boffi F. R., Baron E. 1995, PASP, 107, 1019
Calzetti D., Kinney A. L., Storchi-Bergmann T, 1994, ApJ, 429, 582
Calzetti D., 1997, AJ, 113, 162
Cardelli J.A., Clayton G.C., Mathis J.S., 1989, ApJ 345, 245
Caughlan G. R., Fowler W. A., 1988, At. Data Nucl. Data Tables, 40, 284
Chevalier, R. A. 1976, ApJ 207, 872
Crowther P. A., Dessart L., Hillier D. J., Abbott J. B., Fullerton, A. W., 2002, A&A, 392, 653
Dolphin A.E., 2000a, PASP, 112, 1383
Dolphin A.E., 2000b, PASP, 112, 1397
Dray, L.M., Tout, C. A., Karakas, A. I.,Lattanzio, J.C., MNRAS 2003 in press
Eastman R.G., Schmidt B.P., Kirshner R., 1996, ApJ, 466, 911
Eggleton P. P., 1971, MNRAS, 151, 351
Eggleton P. P., 1972, MNRAS, 156, 361
Eggleton P. P., 1973, MNRAS, 163, 279
Elmhamdi A., et al., 2003, MNRAS, in press, astro-ph/0209623
Evans R., 2001, IAU Circ. 7690
Fassia A., 2001, MNRAS, 325, 907
Ferrarese L., et al., 2000, ApJSS, 128, 431
Filippenko A.V., 1997, ARA&A, 35, 309
Grevesse N., Sauval A.J., 1998, Space Science Rev., 85, 161
Hamuy M., et al, 2001, ApJ, 558, 615
Hamuy M., 2003, ApJ, in press, astro-ph/0209174
Heger A., Woosley S.E., Fryer S.L., Langer N., 2003a, to appear in Proc. of the ESO/MPA/MPE Workshop "From Twilight to Highlight: The Physics of Supernovae", eds. W. Hillebrandt, B. Leibundgut (Springer-Verlag) in press, (astro-ph/0211062)
Heger A., Fryer S.L., Woosley S.E., Langer N., Hartmann D.H., 2003b, ApJ, submitted, astro-ph/0212469
Humphreys R.M., Davidson K., Smith N., 1999, PASP, 111, 1124
Holtzman J.A., et al., 1995, PASP, 107, 156
Holtzman J.A., et al., 1995, PASP, 1065
Iglesias C. A., Rogers F.J., Wilson B.G., 1992, ApJ, 397, 717
Kobulnicky, H.A., Kennicutt, R.C., Pizagno, J.L., 1999, ApJ 514, 544
Krist J., Hook R., 1999, The Tiny Tim Users Guide, wwww.stsci.edu/software/tinytim
Kroupa P., Tout C. A., 1997, MNRAS, 287, 402
Drilling J.S. and Landolt A.U., 2000, in Allen's Astrophysical Quantities, ed. A.N.Cox(4th ed.; New York: AIP)
Lentz E.J., et al., 2001, ApJ, 547, 406
Leonard D.C., et al., 2002a, AJ, 124, 2490
Leonard D.C., et al., 2002b, PASP, 114, 35
Li W., Filippenko A.V., Van Dyk S.D., Hu J., Qiu Y., Modjaz M., Leonard D.C., 2002 PASP, 114, 403
Litvinova I.Y., Nadyozhin D.K., 1985, Sov. Astron. Lett., 11, 145
Mazzali P., et al., 2002, ApJ, 572, L161
McGaugh S.S., 1991, ApJ 380, 140
Meynet G., Maeder A., Schaller G. Schaerer D., Charbonnel C., 1994, A&AS, 103, 97
Nadyozhin D.K., 2003, to appear in Proc. of the ESO/MPA/MPE Workshop "From Twilight to Highlight: The Physics of Supernovae", eds. W. Hillebrandt, B. Leibundgut (Springer-Verlag), in press
Pastorello A., et al., 2002, MNRAS, 333, 27
Pilyugin L.S., 2001, A&A, 369, 584
Pols O.R., Tout C.A., Eggleton P.P., Han Z., 1995, MNRAS 274, 964
Pols O.R., Tout C.A., Schröder K.-P., Eggleton P.P., Manners J., 1997, MNRAS, 289, 869
Pooley D., et al., 2002, ApJ, 572, 932
Popov D.V., 1993, ApJ, 414, 712
Roy J.-R., Walsh J.R., 1997, MNRAS 288, 715
Ryder S., Staveley-Smith L., Dopita M., Petre R., Colbert E., Malin D., Schlegel E., 1993, ApJ, 416, 167
Schaller G., Schaerer D., Meynet G., Maeder A., 1992, A&AS, 96, 269
Schlegel E.M., 2001, ApJ, 556, L25
Silbermann N.A., et al., 1999, ApJ, 515, 1
Smartt S.J., Gilmore G.F., Trentham N., Tout C.A., Frayn C.M., 2001a, ApJ, 556, L29
Smartt S.J., Crowther P.A., Dufton P.L., Lennon D.J., Kudritzki R.P., Herrero A., McCarthy J., Bresolin F., 2001b, MNRAS, 325, 257
Smartt S.J., Kilkenny D., Meikle W.P.S., 2001c, IAUC7704
Smartt S.J., Gilmore G.F., C.A. Tout, S. Hodgkin, 2002a, ApJ, 565, 1089

Smartt S.J., Vreeswijk P. M., Ramirez-Ruiz E., Gilmore G. F., Meikle W. P. S., Ferguson A. M. N., Knapen J. H., 2002b, ApJ, 572, L147

Thompson L.A., 1982, ApJ, 257, L63

Trundle C.A., Dufton P.L., Lennon D.J., Smartt S.J., Urbaneja M., 2002, A&A, 395, 519

Turatto M., Benetti S., Cappellaro E., 2003, to appear in Proc. of the ESO/MPA/MPE Workshop "From Twilight to Highlight: The Physics of Supernovae", eds. W. Hillebrandt, B. Leibundgut (Springer-Verlag), in press, (astro-ph/0211219)

van den Bergh S., Li W., Filippenko A., 2002, PASP, 114, 820

Van Dyk S.D., Hamuy M., Fillipenko A.V., 1996, AJ, 111, 2017

Van Dyk S.D., Peng C Y., King J.Y., Fillipenko A.V., Treffers R.R., Li W., 2000, PASP, 112, 1532

Van Dyk S.D., Li W., Fillipenko A.V., 2003a, PASP, 115, 1

Van Dyk S.D., Li W., Fillipenko A.V., 2003b, PASP, in press, astro-ph/0301346

Vila-Costas M.B., Edmunds M.G., 1992, MNRAS, 259, 121

Walborn N. et al., 1989, A&A, 219, 229

Wang L., Baade D., Fransson C., Hoeflich P., Lundqvist P., Wheeler J.C., 2001, IAU Circ. 7704

White G.L., Malin D.F., 1987, Nat., 327, 36

Whitmore B. et al., 1999, PASP, 111, 1559

Woolley R., Stibbs D. W. N., 1953, The Outer Layers of a Star, Clarendon Press, Oxford

Woosley S.E., Eastman R.G., Weaver T.A., Pinto P.A., 1994, ApJ, 429, 300

Woosley S.E, Weaver T.A., 1986, ARA&A, 24, 205

This paper has been typeset from a TEX/ LATEX file prepared by the author.

This figure "smartt_fig1.jpg" is available in "jpg" format from:

http://arXiv.org/ps/astro-ph/0301324v2